arXiv:1602.02475v1 [math.NT] 8 Feb 2016

Algorithmic construction of Shimura - Taniyama - Weil parametrization of elliptic curves over the rationals

H. Gopalakrishna Gadiyar and R. Padma
Department of Mathematics
School of Advanced Sciences
V.I.T. University, Vellore 632014 INDIA
E-mail: gadiyar@vit.ac.in, rpadma@vit.ac.in

Abstract

In this note we give an algorithm to explicitly construct the modular parametrization of an elliptic curve over the rationals given the Weierstrass function $\wp(z)$.
MSC2010: 11B68, 11F03, 11S40, 14H52, 14L05, 33E05

1 Introduction

We state a straight forward algorithm to construct the modular parametrization of elliptic curves over the rational numbers. The algorithm is implicit in the papers of Honda [5], Clarke [2], Goldfeld [3] and Buchstaber and Bunkova [1] described below. It consists of the following steps:

2 The algorithm

Step 1. Given the equation of the elliptic curve

$$E : y^2 = 4x^3 - g_2x - g_3 \tag{1}$$

over Q, we have the parametrization $x = \wp(z; g_2, g_3), y = \wp'(z; g_2, g_3)$.

Step 2. The formal exponential $f_E(T)$ of the formal group corresponding to this curve has been constructed explicitly by Buchstaber and Bunkova.

$$f_E(T) = -2\frac{\wp(T; g_2, g_3)}{\wp'(T; g_2, g_3)} \tag{2}$$

Step 3. As $f_E(f_L(T)) = f_L(f_E(T)) = T$ the formal logarithm $f_L(T)$ is got from $f_E(T)$ by Lagrange inversion. $f_L(T)$ is related to the L-series of the elliptic curve. This is implicit in the work of Honda and Clarke. If

$$f_L(T) = \sum_{n=1}^{\infty} \frac{a(n)}{n} T^n \tag{3}$$

then

$$L_E(s) = \sum_{n=1}^{\infty} \frac{a(n)}{n^s}\,. \tag{4}$$

and vice-versa.

Step 4. Goldfeld explicitly gives the modular parametrization as

$$x = \alpha(z) = \wp(F(z)) = \wp\left(\sum_{n=1}^{\infty} \frac{a(n)}{n} e^{2\pi i n z}\right) \tag{5}$$

and

$$y = \beta(z) = \wp'(F(z)) = \wp'\left(\sum_{n=1}^{\infty} \frac{a(n)}{n} e^{2\pi i n z}\right)\,. \tag{6}$$

3 Formal group law

A commutative one dimensional formal group law over a ring A is a formal power series ([4], [7])

$$F(t_1, t_2) = t_1 + t_2 + \sum_{i,j \geq 1} a_{ij} t_1^i t_2^j \tag{7}$$

such that the following conditions hold:

$$F(t, 0) = t, F(t_1, t_2) = F(t_2, t_1)$$

and

$$F(t_1, F(t_2, t_3)) = F(F(t_1, t_2), t_3)$$

The formal group F, the formal exponential f_E and the formal logarithm f_L are connected by the relations

$$F(t_1, t_2) = f_E(f_L(t_1) + f_L(t_2)) \tag{8}$$

and

$$f_E(f_L(T)) = f_L(f_E(T)) = T\,. \tag{9}$$

4 Elliptic Curve and its formal group

Let E be an elliptic curve $E : y^2 = 4x^3 - g_2x - g_3$ over Q. Let $\mathcal{L} = \{m\omega_1 + n\omega_2 : m, n \in Z\}$ be the lattice spanned by the two fundamental periods ω_1, ω_2 such that

$$g_2 = g_2(L) = 60 \sum_{l \in \mathcal{L} - \{0\}} \frac{1}{l^4} \tag{10}$$

and

$$g_3 = g_3(L) = 140 \sum_{l \in \mathcal{L} - \{0\}} \frac{1}{l^6}. \tag{11}$$

We have the Weierstrass parametrization $x = \wp(z), y = \wp'(z)$ where $\wp(z)$ the Weierstrass $\wp$ - function

$$\wp(z) = \frac{1}{z^2} + 2 \sum_{n \geq 1} G_{2n+2}(E, \omega) \frac{z^{2n}}{(2n)!} \tag{12}$$

Here

$$G_k = \frac{(-1)^k (k-1)!}{2} \sum_{l \in \mathcal{L} - \{0\}} \frac{1}{l^k} \tag{13}$$

for $k \geq 4$. Note that $G_k = 0$ whenever k is an odd integer. Also all the G_k's can be got in terms of g_2 and g_3.

The Bernoulli -Hurwitz numbers have been defined by Katz[6] as follows. If E is an elliptic curve given by (1), then

$$BH_k = 2kG_k \text{ for } k \geq 4; = 0 \text{ if } k \text{ is odd.} \tag{14}$$

Clarke [2] defines the universal Bernoulli numbers by the formula

$$\frac{T}{f_E(T)} = \sum_{k \geq 0} \hat{B}_k \frac{T^k}{k!}. \tag{15}$$

When $f_E(t) = e^T - 1$, we get the classical Bernoulli numbers B_k.

For the elliptic curve E in the Weierstrass form, Buchstaber and Bunkova [1] have used the theory of solitons to make explicit constructions of formal group laws.

$$F(t_1, t_2) = t_1 + t_2 - b\ m\ \frac{2g_2 + 3g_3 m}{4 - g_2 m^2 - g_3 m^3} \tag{16}$$

is the formal group law of the elliptic curve E, where for $i = 1,\ 2$, $(x_i,\ y_i) \in E$

$$t_i = -2\frac{x_i}{y_i}, s_i = -\frac{2}{y_i}, \tag{17}$$

and

$$m = \frac{s_1 - s_2}{t_1 - t_2},\ b = \frac{t_1 s_2 - t_2 s_1}{t_1 - t_2}\,, \tag{18}$$

and the function

$$f_E(T) = -2\frac{\wp(T; g_2, g_3)}{\wp'(T; g_2, g_3)} \tag{19}$$

is the exponential of the formal group.

Hence the $\hat{B}_k$ can be got in terms of the Bernoulli - Hurwitz numbers BH_k in the case of elliptic curves.

5 Formal logarithm and the L-series of an elliptic curve

The formal logarithm $f_L(T)$ is got from $f_E(T)$ by Lagrange inversion. If

$$f_L(T) = \sum_{n=1}^{\infty} \frac{a(n)}{n} T^n \tag{20}$$

denotes the formal logarithm of an elliptic curve E, then the L- series of the elliptic curve is given by

$$L_E(s) = \sum_{n=1}^{\infty} \frac{a(n)}{n^s}\,. \tag{21}$$

and vice-versa [5].

If we take $f_E(T) = e^T - 1$ in (15), we get the classical Bernoulli numbers B_k and the corresponding formal logarithm is $f_L(T) = \log(1 + T)$ In this case one gets the Dirichlet (L-) series

$$\eta(s) = 1 - \frac{1}{2^s} + \frac{1}{3^s} - \cdots = (1 - 2^{1-s})\ \zeta(s)\,, \tag{22}$$

where $\zeta(s)$ is the classical Riemann - zeta function.

The Shimura - Taniyama - Weil conjecture which is now known as the modularity theorem states that any elliptic curve over rational numbers is modular.

In [3], D. Goldfeld gives the modular parametrization of E as follows:

$$\alpha(z) = \wp(F(z)) = \wp\left(\sum_{n=1}^{\infty} \frac{a(n)}{n} e^{2\pi i n z}\right) \tag{23}$$

and

$$\beta(z) = \wp'(F(z)) = \wp'\left(\sum_{n=1}^{\infty} \frac{a(n)}{n} e^{2\pi i n z}\right) . \tag{24}$$

where

$$f(z) = \sum_{n=1}^{\infty} a(n) e^{2\pi i n z} \tag{25}$$

is a cusp form of weight 2 for $\Gamma_0(N)$ and N is the conductor of the elliptic curve E.

6 Conclusion

The Riemann zeta-function $\zeta(s)$ has two fundamental representations.

$$\sum_{n=1}^{\infty} \frac{1}{n^s} \;=\; \prod_{p} \left(1 - \frac{1}{p^s}\right)^{-1} , \text{ if } Re.\ s \;>\; 1 . \tag{26}$$

These we call the left hand and the right hand side. The left hand side is a sum over the integers and the right hand side is a product over primes. In recent times the p-adic approach which corresponds to the right hand side has been cultivated extensively. It is hoped that the present note will re-kindle an interest in an approach which relies on complex analysis.

7 Acknowledgements

We would like to thank Professors Arul Lakshminarayanan (IIT, Madras), A. Sankaranarayanan (TIFR), K. Srinivas (IMSc, Chennai) for making available the papers and books used in this research.

References

[1] V. M. Buchstaber and E. Yu. Bunkova, Elliptic formal group laws, integral Hirzebruch genera and Krichever genera, arXiv:1010.0944v1

[2] Francis Clarke, The Universal von Staudt theorems, Trans. A. M. S., Volume 115, Number 2, October 1989.

[3] D. Goldfeld, Modular elliptic curves and Diophantine problems, Number Theory, Proceedings of the First Conference of the Canadian Numffber theory Association held at the Banff Centre, Banff, Alberta, April 17-27, 1988, R. Mollin (Ed.), Walter de Gruyter Inc., 157 - 175.

[4] M. Hazewinkel, Formal groups and applications, Academic Press, 1978.

[5] T. Honda, Formal groups and zeta functions, Osaka J. Math. Vol.5, 1968, 199-213.

[6] Nicholas M. Katz, The congruences of Clausen - von Staudt and Kummer for Bernoulli-Hurwitz numbers, Math. Annalen, Vol. 216, 1975, 1-4.

[7] J. Silverman, The arithmetic of elliptic curves, GTM 106, Springer - Verlag, New York, 1985.